

Michigan
Adult Cardiac Protocols
HYPOTHERMIA CARDIAC ARREST

Date: May 31, 2012

Page 1 of 2

Hypothermia Cardiac Arrest

Pre-Medical Control

MFR/EMT/SPECIALIST/PARAMEDIC

1. Follow **General Pre-hospital Care Protocol**.
2. Assess body temperature. If temperature is greater than 30° C (86° F), follow **Cardiac Arrest – General Protocol**
3. If pulseless, begin CPR.
4. Protect against heat loss.
5. Apply heat packs, if available, to axillae, groin, and neck.
6. Administer warmed humidified oxygen, if possible.

SPECIALIST/PARAMEDIC

7. Administer warmed NS IV/IO, if possible.

PARAMEDIC

8. Follow appropriate **VF/VT or Asystole/PEA Protocols EXCEPT**:
 - A. Limit defibrillation to a single attempt.
 - B. Medication Administration: follow appropriate cardiac arrhythmia treatment protocol with 1 round of medication.

EMT/SPECIALIST/PARAMEDIC

9. Initiate transport and contact Medical Control.

Michigan
Adult Cardiac Protocols
HYPOTHERMIA CARDIAC ARREST

Date: May 31, 2012

Page 2 of 2

